

Comparative efficacies of TMR vs RPNI in reducing phantom limb and neuroma pain for patients with BKAs

Annie Cao, Dr. Omri Ayalon

Department of Orthopedics, NYU Langone Orthopedic Hospital

Background

Chronic pain is a common and debilitating complication following lower limb amputation. Neuroma formation and phantom limb pain (PLP) can significantly impact patients' quality of life, with up to 80% of amputees reporting chronic pain. Two surgical innovations, Targeted Muscle Reinnervation (TMR) and Regenerative Peripheral Nerve Interface (RPNI), have emerged as promising techniques for mitigating these outcomes.

This project compares outcomes of TMR and RPNI in below-knee amputees, focusing on phantom limb pain, neuroma pain, and overall pain interference. Patient-reported outcomes were measured using the PROMIS Pain Interference 6b along with surveys on prosthetic use and satisfaction. The goal is to provide early insights into how these techniques impact quality of life, providing guidance in clinical decision-making for surgical treatment of patients undergoing amputations.

TMR	RPNI
Transfer of transected peripheral nerves to motor nerve branches of nearby muscles	Implantation of transected nerve endings into free muscle grafts without motor nerve coaptation
Redirects aberrant nerve signals to prevent pain	Provides a physiologic target for nerve regeneration
Requires available motor nerve branches near amputation site	Requires availability of small free muscle grafts
Potential added benefit for myoelectric prosthetic control	Primarily focused on pain reduction, not prosthetic control

Methods

In the past 7 days...

PAININ3

How much did pain interfere with your enjoyment of life?.....

Not at all

A little bit

Somewhat

Quite a bit

Very much

1

2

3

4

5

PAININ8

How much did pain interfere with your ability to concentrate?.....

1

2

3

4

5

PAININ9

How much did pain interfere with your day to day activities?.....

1

2

3

4

5

PAININ10

How much did pain interfere with your enjoyment of recreational activities?.....

1

2

3

4

5

PAININ14

How much did pain interfere with doing your tasks away from home (e.g., getting groceries, running errands)?.....

1

2

3

4

5

In the past 7 days...

Never

Rarely

Sometimes

Often

Always

PAININ26

How often did pain keep you from socializing with others?.....

1

2


3

4


5

Pain interference was assessed using the **PROMIS Pain Interference Short Form 6b**, a validated patient-reported outcome measure that yields standardized T-scores (mean = 50, SD = 10) benchmarked to the U.S. general population. Additionally, questions involving pain scale, prosthetic use, painkiller use, and overall satisfaction were asked.

Results




PROMIS scores: TMR patients had a mean T-score of **52.4 ± 8.6 (SD)**, with values ranging from 41.0 to 66.4. RPNI patients had a mean T-score of **49.9 ± 8.6 (SD)**, ranging from 41 to 58.1. Both groups scored near the PROMIS population mean of **T = 50**, suggesting that pain interference was not substantially elevated compared to the general U.S. population.



Pain Scale: On a 0–10 scale, average phantom limb pain was **2.28** in TMR patients and **2.0** in RPNI patients, indicating low levels of residual phantom pain in both groups. For neuroma pain, average scores were **1.0** in TMR patients and **2.3** in RPNI patients. Overall, both surgical techniques were associated with low pain ratings, with no clinically meaningful differences between groups.



RPNI procedure done for a large neuroma of the sciatic nerve.



TMR procedure showing peripheral nerves being attached to nearby motor nerves.

Conclusions

1. TMR and RPNI are equally effective at improving pain.
2. TMR and RPNI both restore pain interference levels to normal.
3. TMR and RPNI both average low scores on a scale from 0-10 for neuroma and phantom limb pain
4. 100% of patients reported that they would undergo the procedure again, reflecting very high satisfaction rates.

Future studies should expand the patient population from those who received BKAs to include patients who received AKAs and upper extremity amputations as well. Because limb use and activity level vary by amputation type, outcome levels for TMR and RPNI may differ. In addition, building larger, multicenter cohorts will allow for greater power and more definitive comparisons to guide surgical decision-making.

References

Dumanian, G. A., Potter, B. K., Mioton, L. M., Ko, J. H., Cheesborough, J. E., Souza, J. M., Ertl, W. J., Tintle, S. M., Nanos, G. P., Valerio, I. L., Kuiken, T. A., Apkarian, A. V., Porter, K., & Jordan, S. W. (2019). Targeted Muscle Reinnervation Treats Neuroma and Phantom Pain in Major Limb Amputees: A Randomized Clinical Trial. *Annals of surgery*, 270(2), 238–246. <https://doi.org/10.1097/SLA.0000000000003088>

Hoyt, B. W., Gibson, J. A., Potter, B. K., & Souza, J. M. (2021). Practice Patterns and Pain Outcomes for Targeted Muscle Reinnervation: An Informed Approach to Targeted Muscle Reinnervation Use in the Acute Amputation Setting. *The Journal of bone and joint surgery. American volume*, 103(8), 681–687. <https://doi.org/10.2106/JBJS.20.01005>

Kubiak, C. A., Adidharma, W., Kung, T. A., Kemp, S. W. P., Cederna, P. S., & Vemuri, C. (2022). "Decreasing Postamputation Pain with the Regenerative Peripheral Nerve Interface (RPNI)". *Annals of vascular surgery*, 79, 421–426. <https://doi.org/10.1016/j.avsg.2021.08.014>

Lin Z, Yu P, Chen Z, Li G. Regenerative peripheral nerve interface reduces the incidence of neuroma in the lower limbs after amputation: a retrospective study based on ultrasound. *J Orthop Surg Res*. 2023;18(1):619. Published 2023 Aug 24. doi:10.1186/s13018-023-04116-6

Senger JB, Hardy P, Thorkelsson A, et al. A Direct Comparison of Targeted Muscle Reinnervation and Regenerative Peripheral Nerve Interfaces to Prevent Neuroma Pain. *Neurosurgery*. 2023;93(5):1180-1191. doi:10.1227/neu.0000000000002541

Yuan M, Gallo M, Gallo L, et al. Targeted Muscle Reinnervation and Regenerative Peripheral Nerve Interfaces Versus Standard Management in the Treatment of Limb Amputation: A Systematic Review and Meta-Analysis. *Plast Surg (Oakv)*. 2024;32(2):253-264. doi:10.1177/22925503221107462

Acknowledgements

This project is a result of the NYU Langone Orthopedic Hospital Surgery Summer Externship Program and completed under the guidance of Dr. Omri Ayalon. Thank you as well to Dr. Steven Glickel and Gabriele Lippert for their support and contributions!